



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE GENUS CRYPTOTOMUS COPE

A recent examination of the type species (*C. roseus* Cope), compared with other members of this genus, convinces me that at least two well-marked groups occur, which may be admitted to sub-generic rank. They may be set forth as follows:

Sub-genus *Cryptotomus* Cope.

Type *Cryptotomus roseus* Cope.

Body elongate, labroid in appearance. Front profiles attenuate. Cutting-edge of jaw teeth formed of several overlapping series, each tooth distinct, and posterior flaring out behind. Scales on breast greatly enlarged. One row of large scales on cheek. Dorsal spines very flexible.

One species, brightly colored.

Sub-genus *Nicholsina* sub-gen. nov.

Type *Cryptotomus beryllinus* Jordan and Swain.

Body moderately long, scaroid in appearance. Front profiles gently convex. Cutting-edge of jaw teeth formed of nearly continuous or single series, with only few anterior distinct. Free posterior canines frequent. Scales mostly uniform in size, not enlarged on breast. Cheek with one or two rows of large scales. Dorsal spines flexible.

Several species, small, and usually of dull or olive coloration. Besides the type, they are *C. dentiens* (Poey), *C. retractus* (Poey), *C. ustus* (Valenciennes) and *C. auropunctatus* (Valenciennes), in America.

Dedicated to Mr. John T. Nichols, of the American Museum of Natural History.

HENRY W. FOWLER,
Philadelphia, Pa.

A NOTE ON A DRAWING OF CARANX BARTHOLOMAEI CUVIER & VALEN- CIENNES.

In the Fishes of North and Middle America, Jordan & Evermann (Bull. 47, U. S. Nat. Mus. vol. iv., 1900), plate CXLII, fig. 389 is erroneously labelled *Caranx latus*. This drawing was made by H. L. Todd from a specimen collected by Vinal N. Ed-